

Frequency dependent transport in the integer quantum Hall effect

A. Bäker, L. Schweitzer

Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany

Abstract The frequency dependent transport is investigated for a two-dimensional disordered system under QHE conditions. The real and imaginary parts of the conductivity are calculated numerically in linear response using a recursive Green function technique. The energy dependence of $\sigma_{xx}(E, \omega)$ is obtained within the lowest Landau band. When the width of the system exceeds the characteristic length, $L_\omega = (\hbar\omega\rho(E))^{-1/2}$, the maximum of the real part of $\sigma_{xx}(E, \omega)$ decays with frequency almost linearly which is different from the classical Drude behaviour.

1 Introduction

The integer quantum Hall effect is usually explained by particular localisation-delocalisation effects that emerge when electrons move in disordered two-dimensional systems in the presence of a strong perpendicular magnetic field. These phenomena can be investigated in more detail if one considers dynamic processes such as frequency dependent electrical transport. Although electron-electron interactions are commonly neglected, because they are believed to be of minor importance in the integer QHE, analytical theories that allow the calculation of the transport coefficients are not available nowadays. Therefore, numerical investigations are a practical means for computing the desired quantities.

In previous numerical work, low frequency anomalies and size scaling of the real part of the dynamic conductivity have been reported [1,2]. Recently, we were able to calculate the frequency dependence of the real and imaginary part of $\sigma_{xx}(\omega)$ in the lowest Landau tail [3] which could be compared with approximate analytical predictions [4]. Also, the frequency scaling of $\Re\sigma_{xx}(\omega)$ in the lowest Landau band has been discovered where a power law $\sim \omega^\kappa$ was observed with $\kappa = (z\mu)^{-1} \simeq 0.2$ [3] which leads to a critical exponent $\mu \approx 2.35$ [5] with an assumed $z = 2$ for non-interacting electrons. This is in good agreement with the frequency scaling results of Engel *et al.* [6] (except that $z = 1$ due to interactions), but at variance with the conclusions of Ref. [7]. An interpretation of the conflicting experimental findings is given in [8]. In the present contribution we focus on both the real and imaginary parts of the frequency dependent longitudinal conductivity $\sigma_{xx}(\omega)$ in the centre, and on the energy dependence within the lowest Landau band.

2 Model and Method

We consider a disordered two-dimensional electronic system in a strong magnetic field with z -component B described by a tight binding Hamiltonian. The magnetic field, which is expressed by a vector potential in Lan-

dau gauge, $\mathbf{A} = (0, Bx, 0)$, is introduced via complex phase factors $V_{(x_m, y_m); (x_m, y_m \pm a)} \sim \exp(\pm i2\pi\alpha_B x_m/a)$ in the transfer terms for transitions between neighbouring lattice sites along the y -direction [3,9]. We choose $\alpha_B = eBa^2/h = 1/8$ which corresponds to $1/8$ flux quantum per lattice cell a^2 . The uncorrelated disorder potentials are distributed uniformly between $-W/2$ and $W/2$. The conductivity is calculated within linear response from the Kubo-formula iteratively by a generalisation of a recursive Green function technique introduced earlier [10,11]. For the frequency dependent longitudinal conductivity at zero temperature we get

$$\begin{aligned} \sigma_{xx}(\omega) = \lim_{\varepsilon \rightarrow 0} \lim_{\Omega \rightarrow \infty} \frac{e^2/h}{\Omega\hbar\omega} & \left[\int_{E_F - \hbar\omega}^{E_F} dE \operatorname{Tr} \left[(\hbar\eta)^2 x G_\omega^+ x G^- \right. \right. \\ & - (\hbar\omega)^2 x G_\omega^+ x G^+ + 2i\varepsilon x^2 (G_\omega^+ - G^-) \\ & \left. \left. - \int_{-\infty}^{E_F - \hbar\omega} dE \operatorname{Tr} \left[(\hbar\omega)^2 \{ x G_\omega^+ x G^+ - x G_\omega^- x G^- \} \right] \right] \right], \quad (1) \end{aligned}$$

where $\Omega = LM$ is the area of the system, $\eta = \omega + 2i\varepsilon/\hbar$, and $G_\omega^\pm = (E - H + \hbar\omega \pm i\varepsilon)^{-1}$ are the advanced and retarded Green function, respectively.

3 Results and Discussion

3.1 Frequency dependence at filling factor $\nu = 1/2$

The results for the real part of the peak value of the longitudinal conductivity $\sigma_{xx}(\omega)$ versus frequency ω are shown in Fig. 1 for two different values of disorder. For small frequencies an almost linear decay is observed for disorder strengths $W/V = 0.1$ and 1.0 . This is in contrast to the classical Drude behaviour given by

$$\sigma_D(\omega) = \sigma_0 \frac{1 + \tau^2(\omega_c^2 + \omega^2)}{1 + 2\tau^2(\omega_c^2 + \omega^2) + \tau^4(\omega_c^2 - \omega^2)^2}, \quad (2)$$

where $\omega_c = eB/m$ is the cyclotron frequency and τ is the collision time. In the limit $\omega \ll \omega_c$, the Drude conductivity $\sigma_D(\omega)$ is almost constant whereas the peak value of $\Re\sigma_{xx}(\omega)$ clearly decays with frequency. A similar non-Drude decay has been reported in Ref. [2] for a different type of model with strong correlated disorder potentials. Owing to the knowledge of the results for a semiclassical model [12] the deviations from Drude-like behaviour have been attributed [2] to long time tails in the velocity correlations which were shown to exist in a QHE system in the absence of electron-electron interactions. The effects of interactions have been taken into account in Ref. [13].




Fig. 1 The real part of the longitudinal conductivity $\sigma_{xx}(\omega)$ in the centre of the lowest Landau band as a function of frequency ω for disorder strengths $W/V = 0.1$ (+), and $W/V = 1.0$ (x). The decay of the data, which are shown in comparison with linear relations (broken lines), is in contrast to the classical Drude behaviour. The width and length of the systems are $M/a = 32$ and $L/a = 2 \cdot 10^6$, respectively.

For the tight-binding model with uncorrelated disorder potentials considered in the present paper it is not clear whether the picture of electron motion along equipotential lines, a basic ingredient for the arguments of Ref. [2], is appropriate. Nevertheless, similar non-Drude behaviour is seen in the frequency dependence of the peak value of the real part of $\sigma_{xx}(\omega)$ in the lowest Landau band. We notice that the frequency decay of $\sigma_{xx}(\omega)$ is accompanied by the decrease of the characteristic length $L_\omega = (\hbar\omega\rho(E))^{-1/2}$ below the system width M . Recently, an increase of $\Re\sigma_{xx}(\omega)$ with enhancing frequency at the QHE transition has been reported for a model based on Dirac fermions [14].

The frequency dependence of the imaginary part of $\sigma_{xx}(\omega)$ for filling factor near $\nu = 1/2$ is shown in Fig. 2. As before, the disorder strength and system width are $W/V = 0.1$ and $M/a = 32$, respectively. For small frequencies a behaviour $\Im\sigma_{xx}(\omega) \sim \omega$ is observed. The linear increase tapers off at about $\hbar\omega/V = 0.015$ and saturates at a value of $\Im\sigma_{xx}(\omega) \simeq 2.39$.




Fig. 2 The imaginary part of the longitudinal conductivity $\sigma_{xx}(\omega)$ in the centre of the lowest Landau band as a function of frequency ω for disorder strength $W/V = 0.1$. The width and length of the system is $M/a = 32$ and $L/a = 2 \cdot 10^6$.




Fig. 3 The energy dependence of the real ($\diamond$) and imaginary (+) part of the longitudinal conductivity $\sigma_{xx}(\omega, E)$ in the lowest Landau band. The applied frequency is $\hbar\omega/V = 0.001$ and the disorder strength $W/V = 0.1$. The width and length of the system is $M/a = 32$ and $L/a = 2 \cdot 10^6$, respectively.

3.2 Energy dependence of $\sigma_{xx}(\omega)$

The energy dependence of both the real and imaginary parts of $\sigma_{xx}(\omega)$ are shown in Fig. 3 for frequency $\hbar\omega/V = 0.001$, and disorder strength $W/V = 0.1$. The system width is $M/a = 32$ so that L_ω is smaller than M . While $\Re\sigma_{xx}(\omega)$ shows the expected conductivity peak the imaginary part exhibits a weak positive double peak structure before it becomes negative at the upper Landau band edge. We note here that the behaviour of $\Im\sigma_{xx}(\omega)$ changes completely, i.e., $\Im\sigma_{xx}(\omega)$ takes negative values for almost any energy, if $L_\omega > M$. Further numerical and analytical work is necessary to elucidate the microscopic origin of the non-Drude behaviour of the peak value of $\Re\sigma_{xx}(\omega)$ and the peculiar frequency dependence of the imaginary part.

References

1. B. M. Gammel and W. Brenig, Phys. Rev. Lett. **73**, (1994) 3286.
2. B. M. Gammel and W. Brenig, Phys. Rev. B **53**, (1996) R13279.
3. A. Bäker and L. Schweitzer, Ann. Phys. (Leipzig) **8**, (1999) SI-21.
4. O. Viehweger and K. B. Efetov, J. Phys.: Condens. Matter **2**, (1990) 7049.
5. B. Huckestein, Europhysics Letters **20**, (1992) 451.
6. L. W. Engel, D. Shahar, C. Kurdak, and D. C. Tsui, Phys. Rev. Lett. **71**, (1993) 2638.
7. N. Q. Balaban, U. Meirav, and I. Bar-Joseph, Phys. Rev. Lett. **81**, (1998) 4967.
8. F. Kuchar, R. Meisels, and B. Kramer, Advances in Solid State Physics **39**, (1999) 231.
9. L. Schweitzer, B. Kramer, and A. MacKinnon, J. Phys. C **17**, (1984) 4111.
10. A. MacKinnon, Z. Phys. B **59**, (1985) 385.
11. L. Schweitzer, B. Kramer, and A. MacKinnon, Z. Phys. B **59**, (1985) 379.
12. F. Evers and W. Brenig, Z. Phys. B **94**, (1994) 155.
13. W. Brenig, B. M. Gammel, and P. Kratzer, Z. Phys. B **103**, (1997) 417.
14. G. Jug and K. Ziegler, Phys. Rev. B **59**, (1999) 5738.